



Recruitment of oysters of *Crassostrea* genus (Bivalvia, Ostreidae) in Guaratuba Bay, PR, Brazil

SUSETE W. CHRISTO^{1*}, AUGUSTO L. FERREIRA-JR^{1,2}, ANDRÉA C. CRUZ-KALED² & THERESINHA M. ABSHER²

¹ Universidade Estadual de Ponta Grossa (UEPG), Departamento de Biologia Geral. Av. Carlos Cavalcante, 4748 – Uvaranas – CEP: 84030-900 – Ponta Grossa – Paraná - Brasil.

² Universidade Federal do Paraná (UFPR), Centro de Estudos do Mar. Av. Beira Mar, s/n - Pontal do Sul – CEP: 83255-000 – Pontal do Paraná – PR – Brasil.

*Corresponding author: swchristo@hotmail.com

Abstract. Larval development of *Crassostrea* is characterized by three larval phases: larva D, Umbo and Pediveliger. In Pediveliger the larvae are near settlement and metamorphosis in hard substratum. In this work, recruitment of oysters' larvae of genus *Crassostrea* (*C. rhizophorae* and *C. brasiliana*) was studied in Guaratuba Bay (Paraná – Brazil) through analysis of artificial collectors placed in three points: Point I (bay's entrance), Point II (cultivation park) and Point III (medium sector of the bay) and changed biweekly from June/2003 to June/2004. Due to the impossibility to differentiate the recruits of the two oysters' species by the shell morphology, the oyster's spats of the two species were counted jointly on both sides of the plates, to determine the number of settled individuals. The results showed an increase in the number of settled individuals between October and March with successive peaks during this period. The medium sector of the bay showed the larger density of recruits along the studied period. An important application of the present research would be in the improvement of oyster cultivation based in seed collection from natural environment.

Key-words: larvae, collectors, spats, Guaratuba Bay, South Atlantic.

Resumo. Recrutamento de ostras do gênero *Crassostrea* (Bivalvia, Ostreidae) na baía de Guaratuba, PR, Brasil. O desenvolvimento das larvas de ostras do gênero *Crassostrea* é caracterizado em fases larvais: larva D, umbo e Pediveliger. No estágio Pediveliger as larvas estão próxima da metamorfose e assentamento no substrato consolidado. Neste trabalho o recrutamento das larvas de ostras do gênero *Crassostrea* (*C. rhizophorae* e *C. brasiliana*) foram estudados na baía de Guaratuba (Paraná, Brasil) através da análise de coletores artificiais em três locais: ponto I (entrada da baía), ponto II (parque de cultivo) e ponto III (setor médio da baía) e a variação quinzenal entre junho/2003 a junho/2004. Devido à impossibilidade de diferir os recrutas das duas espécies nativas de ostras a partir da morfologia da concha, as sementes das duas espécies foram contadas em conjunto de ambos os lados das placas, para determinar o número de indivíduos fixados. Os resultados mostraram um aumento no número de indivíduos fixados entre os meses de outubro e março, com sucessivos picos durante este período. O setor médio da baía mostrou a maior densidade de recrutas ao longo do período estudado. Uma aplicação importante da presente pesquisa seria no aperfeiçoamento do cultivo de ostras com base na coleta de sementes do ambiente natural.

Palavras chaves: larvas, coletores, sementes, baía de Guaratuba, Atlântico Sul.

Introduction

During early developmental stages (D-larvae

and Umbo-larvae), *Crassostrea* spp. larvae are abundant in surface plankton, sinking to deeper

waters during late (pediveliger) stages (Andrews 1983, Boehs & Absher 1997, Baker 2003, Finelli & Wethey 2003). At settlement, substrate selection is a key factor controlling species distribution at local scales (Southworth & Mann 2004, Christo & Cruz 2009). The recruitment process in space and time scales can also supply information about the behavior of the adult oyster in the environment through the observation of the patterns of reproduction, spawns, estimates of the larval period, dispersion and growth associated to environmental factors (Southworth & Mann 2004, Wilson *et al.* 2005). Among these factors temperature, salinity, primary productivity and local hydrodynamic are very important (Boehs & Absher, 1997, Nalesso *et al.* 2008, Kim *et al.* 2010, Haase *et al.* 2012).

Researches with artificial substrate are quite spread in oyster culture, and have been used especially in areas where the artificial production of larvae in laboratory is unviable due to the high cost or when the objective is to increase settlement rate in the environment. In the recruitment process the interaction among genetic and environmental factors can also contribute to the substratum selection and settlement rate (Baker 2003, Finelli & Wethey 2003, Silveira *et al.* 2011). In the present study, the selection of the polypropylene substratum was based on previous works accomplished in Paranaguá Bay, on the influence of the substratum in the settlement of oysters' larvae (Absher & Christo 1991, Christo & Absher 1993). The selection of the type of artificial substratum for the collectors is important in the optimization of spat recruitment in the nature (Sorabella & Luckenbach 2003, Jordan & Coakley 2004, Padilla & Klinger 2005, Su *et al.* 2007). The emphasis of the studies has been in the selection of the size of the collectors and materials of low cost and handling with the optimization purpose in the recruitment of larvae in nature (Mann 2001, Sorabella & Luckenbach 2003, Jordan & Coakley 2004, Padilla & Klinger 2005, Christo & Cruz 2009, Zanette *et al.* 2009). Therefore, this work had as objective to verify the space and time variation of larval recruitment of *Crassostrea* oysters (*C. rhizophorae* and *C. brasiliana/C. gasar*) through artificial collectors and identify their relationship to the environmental variables that act on the process of settlement of the seeds in the natural environment.

Materials and Methods

Guaratuba Bay situated on the coastal plain of southern Brazil (Fig. 1), has an area of 50.19 km² and four cross-estuary sections (bay mouth; middle;

São João river mouth; Cubatão river mouth) were established (Marone *et al.* 2006). The climate of the region is humid mesothermal. Marone *et al.* (2006) describes very dynamic situation, with strong currents (more than 2.0 m.s⁻¹ at the mouth), semi-diurnal tides of up to 1.5 m range (0.65 m at neap tide), moving up to 75x10⁶m³ of water at spring tide and 32x10⁶km² at neap tides in a complex tide propagation pattern (progressive, mixed and stationary).

Oyster's recruitment was analyzed, in the period from June 2003 to June of 2004, through artificial collectors placed at three locals of Guaratuba Bay, as follows: Point I (25°51'20"S; 48°34'39"W – bay mouth) - close to Sepultura's Island, in the entrance of the bay, where natural banks of oysters occur in rocky coasts. Point II (25°49'56"S; 48°34'48"W - middle) - is located in a cultivation park of *C. brasiliana* in Pinheiros' River, formed by a tide channel in the northeast portion of the bay between Veiga Island and the continent. This channel has an approximate width of 150m, length of 2000m. Point III (25°52'12"S; 48°38'21"W - middle) - near Garças' Island, in the medium sector of the bay has a mean depth of (Fig. 1). In each point 2 collectors with 5 plates of polypropylene each were placed, to 1 m depth from the sea surface. This depth was selected in order to prevent the establishment of native oyster according with preference of larvae settlement. *C. rhizophorae* is predominant intertidal and *C. brasiliana/C. gasar* is predominant infralittoral. A total of 10 plates were disposed in strings of galvanized wire and each plate with an area of 10 cm² was placed 4 cm apart. Every 15 days the collectors were substituted and taken to the laboratory. Jointly to spat sampling environmental factors were measured: temperature with a mercury thermometer - precision of 0.1 °C; salinity with manual refractometer - precision of 1 unit; seawater transparency with Secchi disk. Pluviosity data was obtained from the meteorological station of Guaratuba of the Technological Institute SIMEPAR (Meteorological System of Paraná).

Due to the impossibility to differentiate the recruits of the two oysters' species by the shell morphology, the oyster's spats of the two species were counted jointly on both sides of the plates, to determine the number of settled individuals. The number of spats settled in each plate was transformed in densities, expressed in individuals/100 cm².




Figure 1 - Location of points in Guaratuba Bay, Paraná – Brazil. Points: I (near Sepultura Island), II (cultivation park) and III (medium sector of the bay). Sections: bay mouth; middle; São João river mouth; Cubatão river mouth (Marone *et al.* 2006).

Spats data were transformed (square root). After evaluated the normality and homogeneity of the data, one-way ANOVA was used to test for significant differences ($\alpha=0.05$) in larvae densities (dependent variable) between plates (1 to 5) and side of the plate area (upper and lower). A two-way ANOVA was used to test for significant differences ($\alpha=0.05$) in larvae densities (dependent variable) between points and stations sampled (independent variable). The correlation between the biological parameters and the environmental parameters was put in evidence through a multivariate analysis of ordination as a PCA (Principal Component Analysis) (Legendre & Legendre 1983, Valentin 2000), with the seasonal averages of the variables.

Results

Recruitment of oysters: The analysis of the results of the recruitment of oysters showed an increase in the number of individuals settled between the months of October/2003 and March/2004 with successive picks during this period. An annual average of 112 ± 77.61 spats.100cm⁻² was observed at Point I; Ponto II, 81 ± 59.59 spats.100cm⁻² and Ponto III an average of 193 ± 22.71 spats.100cm⁻². The largest densities observed were in spring (October/2003) and autumn (March/2004), where the Ponto III (medium section of the bay) showed a larger density of settled

individuals (679.5 spats.100cm⁻²) in this period (Fig. 2).

Comparing the sampled points, the variance analysis showed that the differences in the recruits' densities were highly significant ($F=49.35$; $p<0.001$) (Fig. 3). There was significant difference between the number of recruits and the period sampled ($F=68.78$; $p=0.001$) (Fig. 4).

In the three studied points a more expressive recruitment is observed at springtime and a lower amount in the winter period ($F=14.20$; $p<0.01$) probably justifying the high significant interaction between the season and recruits (Fig. 2-4). Comparisons of density values of spats among plates ($p=0.9183$) and the upper and lower inferior side of the plate ($p=0.2559$).

Environmental parameters: Seawater temperature had an annual variation of 7.4°C with values that oscillated between 19°C and 27°C and annual average of 23°C in Points I, II and III. The highest temperatures were registered in the period from spring (October/2003) to autumn (March/2004). The annual average of the salinity was 26 in the Ponto I; 24 in Point II and 19 in Point III, with annual amplitude of 10, 6 and 13 in the Points I, II and III, respectively. The smallest values were registered in Point III, at the medium area of the bay.




Figure 2 - Monthly average of the number of recruits in Points I (light grey), II (dark grey) and III (white) during the studied period in Guaratuba Bay, Paraná – Brazil.

The annual average of the depth of extinction of the Secchi Disk was 1.60 m in the Ponto I; 1.15 m in Point II and 1.25 m in Point III, with annual amplitude of 1.10, 0.80 and 1.90 m in the Points I, II and III, respectively. The largest value of transparency of the water was verified in the Point I (2.20 m) that corresponds to the area at the entrance of the bay and the smallest registered value (0.50 m) was registered in summer (February/2004) in Point III.




Figure 3 - Distribution of the number of recruits (spats.100cm⁻²) in Points I, II and III, during the studied period in Guaratuba Bay, Paraná – Brazil. ●: mean; C: stander error; ▭: stander deviation.

Environmental and biological interactions: Result of

PCA analysis evidenced that in the points I, II and III the recruits' density was accentuated in the spring/2003 and summer/2004 in conditions of high temperature and less saline waters. Pluviosity had little influence on the recruits' density, while water transparency didn't interfere in the recruitment. In Points I and III an inverse relationship of recruitment to the transparency of the water was observed, however, the salinity didn't have influence on the recruits' density (Fig. 5).

Discussion

In this study, the settlement of oysters' spat, does not present a preference close to the bottom, as verified by Wakamatsu (1973), Akaboshi & Pereira (1981), Christo & Cruz (2009) for these organisms; because significant differences were not observed in the number of recruits between the upper and lower sides of the plates. This behavior can be related to the little depth of the bay and/or temperature, since Andrews (1983) suggests that oysters' larvae tend to stay in layers of water masses with more favorable temperature. Nalesso *et al.* (2008) observed spat settlement was significantly higher on oyster shell, tile and tire collectors. Hossain *et al.* (2013) in three sites in the Bangladesh coast observed higher performance of windowpane shell for spat settlement when comparing four different substrates: dead oyster shell, live oysters, windowpane shell and stone/rubble. Silveira *et al.* (2011) in laboratory

experiments of larval settlement and spat recovery rates of *Crassostrea brasiliensis* selected plastic collectors due to its pliability, favoring spat detachment.

A continuous reproductive pattern is observed in the oyster from Guaratuba Bay what favors the presence of larvae in the plankton along the whole year with temperature playing an important role in this pattern, high density of larvae and gonadal maturation coincides with periods of temperature elevation (Christo 2006). This fact indicates that the spawning activity influence the pattern of recruitment of the oysters (Southworth & Mann 2004, Wilson *et al.* 2005). Nalesso *et al.* (2008) in five sites in the Benevente river estuary, Espírito Santo State – Brazil observed that recruitment of oysters' spat occurred throughout the year, being higher from November to February. Similar increase was observed in recruitment of oysters' spat of Guaratuba bay, where there was an increase from October to March.

Related with water transparency the highest values correspond to periods where the rate of

rainfall was low, especially verified in the winter months; while in the hottest and rainy periods there was an accentuated increase in turbidity. This fact might have happened due to the development of phytoplankton and/or suspension of the sediment due to little depth of the area, especially verified in Point II that corresponds to the medium sector of the bay. The river basins that contribute for the total runoff into the Guaratuba bay represents a surface area of 1,724 km², with a drainage density of 1.87 rivers.km⁻¹, presenting a dendritic drainage pattern and a mean river runoff of at least 80m³.s⁻¹ (Marone *et al.* 2006). It is important to note, also, the role of the relative importance between tides and river runoff, which seasonal changes, moving the bay's dynamic from a highly stratified to a well mixed estuary in different time scales (Marone *et al.* 2006).

These patterns of surface sediment being dragged in the direction of the inside of the bodies of water, mainly during the periods of intense rains, are according with the results of the transparency of the water obtained in the present study.




Figure 4 - Seasonal averages of the number of recruits (spats.100cm⁻²) in Points I (○), II (□), and III (Δ) during the studied period in Guaratuba Bay, Paraná – Brazil. $\pm$: standar deviation.




Figure 5 - Graphic representation of Principal Components Analysis (PCA) in Points I, II and III, with the projection of the vector-variables, being: R = recruits; T = temperature ($^{\circ}\text{C}$); S = salinity; Tb = water transparency (m) and Plu = pluviosity (mm); Sum = summer; Aut = autumn; Win = winter; Sprin = spring, with the Seasonal observations in the factorial plan I and II. Mean seasonal values in the sampled period from June/03 to June/04.

There is a conspicuous stratification attributed to the geomorphologic characteristics of the Guaratuba bay with an entrance channel strait and deep, added by the time of the water residence calculated by the referred authors as being of approximately 9.3 days (Marone *et al.* 2006), this characteristic may contribute to the retention of the larvae in the internal sections of the bay. Study to investigate the larval transport of eastern oyster, *Crassostrea virginica*, in two American estuarine systems indicated the influence of hydrodynamics, seasonality and the population dynamics of oysters increased larval retention near the spawning area, thus providing a favorable condition for local recruitment of oysters (Kim *et al.* 2010, Haase *et al.* 2012).

However, the analysis of the environmental parameters, suggests that the hottest waters favor the recruitment and the permanence of larvae in the plankton, while no direct relationship was observed in relation to the water transparency in the studied points. Finally, those results may have an important application in commercial exploitation of oyster farms in the area that depends on seed collection from natural environment. Nalesso *et al.* (2008) recommend that collectors be placed in the water from October to December (to collect spat during the rainy months) in high salinity areas, and that they should be transferred to areas of low salinity

from April, to avoid the rainy season at estuarine sites. An accentuated reduction in larvae and recruits density was observed in February/2004 in Point III, where the smallest value of water transparency (0.50 m) was registered. This fact can be justified by the period of the collectors' substitution. Point III, close to Garças' Island, is a place often visited by tourists' boats of small and medium size especially in this time of the year. The great circulation of high-speed boats in this area can cause the sediment suspension and displacement of water, what probably interfered with the distribution of larvae and recruits due to disturbances caused in the environment in this period. In this estuarine system 8 marina can be identified located in the cities of Matinhos and Guaratuba (SETPT 2012), as well as continuous flow of ferry boats in their outlet, responsible for the crossing Matinhos - Guaratuba that happens daily along the year (DER 2015). According with DER (2015), in the summer period 3 ferry-boats and 3 tug boats are used for 24 hours to assist the demand of the tourist season, besides the presence of a shipyard of maintenance of the six embarkations (ferry-boats and tug boats).

Analyzing evidencing the recruitment seasonally, being the seasons the independent variables, Point I showed a high number of individuals in autumn, in the Point II in spring and Ponto III in autumn and spring, this fact could be associated to the elevation of the seawater temperature, salinity and water transparency. Perbiche-Neves *et al.* (2010) in five sites in Guaratuba Bay (sections: bay mouth and middle) observed spatial and seasonal variations for the micro-zooplankton in the bay, with peaks of abundance (due to meroplankton) in the summer, associated to the highest water temperatures. The number of oyster spat was positively correlated with the salinity and water temperature, revealing that areas with higher salinities and summer months were better for spat collection in the Benevente river estuary, Espírito Santo State - Brazil (Nalesso *et al.* 2008). Studies observed the number of oyster settlements in three locations of Bangladesh coast indicated that main factors were an increase in suspended solids, low salinity, due to heavy rains, increased river discharge by months of monsoon high water flow rates and current wave in extreme weather events. All those factors can affect the success of the spats survival in the experimental sites (Hossain *et al.* 2013).

References

- Absher, T. M. & Christo, S. W. 1991. Efeito do material do substrato artificial no recrutamento de ostras e cirripédios. **SBMa XII Encontro Brasileiro de Malacologia, São Paulo. Sociedade Brasileira de Malacologia-SBMa**, pp.45. Accessible at http://www.sbma.uerj.br/_recursos/pages/ebram_1991.php Accessed 15 March 2016.
- Akaboshi, S. & Pereira, O. M. 1981. Ostreicultura na região lagunar de Cananéia, São Paulo, Brasil; 1. Captação de larvas da ostra *Crassostrea brasiliiana* (Lamarck, 1819) em ambientes natural. **Boletim Instituto de Pesca**. 8: 87-104.
- Andrews, J. D. 1983. Transport of bivalve larvae in James River, Virginia. **Journal of Shellfish Research** 3: 29-40.
- Baker, P. 2003. Two species of oyster larvae show different depth distributions in a shallow, well-mixed estuary. **Journal of Shellfish Research** 3: 733-736.
- Boehs, G. & Absher, T. M. 1997. Distribuição de larvas de ostras do gênero *Crassostrea* SACCO, 1897 (OSTREOIDA:OSTREIDAE) na Baía de Paranaguá, Paraná. **Arquivos de Biologia e Tecnologia**. 40: 39-45.
- Bushek, O. 1988. Settlement as a major determinant of intertidal oysters and barnacle distributions along a horizontal gradient. **J. Exp. Marine Biology and Ecology**. 122: 1-18.
- Christo, S. W. & Absher, T. M. 1993. Recrutamento de ostras do gênero *Crassostrea* SACCO, 1897 na Gamboa Perequê – Baía de Paranaguá, PR. **SBMa XIII Encontro Brasileiro de Malacologia, Rio de Janeiro. Sociedade Brasileira de Malacologia-SBMa**, pp. 56. Accessible at http://www.sbma.uerj.br/_recursos/pages/ebram_1993.php. Accessed 15 March 2016.
- Christo, S. W. 2006. Biologia reprodutiva e ecologia de ostras do gênero *Crassostrea* Sacco, 1897 na Baía de Guaratuba (Paraná – Brasil): um subsídio ao cultivo. **Tese de doutorado**. Universidade Federal do Paraná, Departamento de Zoologia, 137pp.
- Christo, S. W. & Cruz, E. 2009. Recrutamento de ostras e cracas (Cirripédia) em um canal artificial de Pontal do Sul, Pontal do Paraná, PR. **Publication**. 15(2): 43-48.
- DER. 2015. **Departamento de Estradas de Rodagem - FERRY BOAT-DER**, accessible at <http://www.der.pr.gov.br/modules/conteudo/conteudo.php?conteudo=74> 1-2p. Accessed 17 August 2015.
- Finelli, C. M. & Wetthey, D. S. 2003. Behavior of oyster (*Crassostrea virginica*) larvae in flume boundary layer flows. **Marine Biology**. 143: 703-711.
- Haase, A. T., Eggleston, D. B., Luettich, R. A., Weaver, R. J., & Puckett, B. J. 2012. Estuarine circulation and predicted oyster larval dispersal among a network of reserves. **Estuarine, Coastal and Shelf Science**. 101, 33-43.
- Hossain, M. S., Rothuis, A., Chowdhury, S. R., Smaal, A., Ysebaert, T., Sharifuzzaman, S. M., van Sluis, C., Hellegers, P., van Duijn, A., Dankers, P. & Chowdhury, S. N. 2013. Oyster aquaculture for coastal defense with food production in Bangladesh. **Aquaculture Asia**. 18(1): 15-24.
- Jordan, S. J. & Coakley, J. M. 2004. Long-term projections of eastern oyster populations under various management scenarios. **Journal of Shellfish Research**. 23: 63-72.
- Kim, C. K., Park, K., Powers, S. P., Graham, W. M., & Bayha, K. M. 2010. Oyster larval transport in coastal Alabama: Dominance of physical transport over biological behavior in a shallow estuary. **Journal of Geophysical Research: Oceans**. 115(C10019), 1-16.
- Legendre, L. & Legendre, P. 1983. **Numerical ecology Developments in Environmental Modelling**. Elsevier Scientific Publishing Company, Amsterdam. 412pp.
- Mann, R. 2001. Restoration of the oyster resource in Chesapeake Bay: The role of oyster reefs in population enhancement, water quality improvement and support of diverse species-rich communities. **Bulletin of the Aquaculture Association of Canada**. 101: 38-42.
- Marone, E., Noernberg, M. A., Santos, I., Lautert, L. F. C., Andreoli, O., Buba, H. & Fill, H. D. 2006. Hydrodynamic of Guaratuba bay PR, Brazil. **Journal Coastal Research**. 39: 1879-1883.
- Nalesso, R. C., Paresque, K., Piumbini, P. P., Tonini, J. F. R., Almeida, L. G., & Nickel, V. M. (2008). Oyster spat recruitment in Espírito Santo State, Brazil, using recycled materials. **Brazilian Journal of Oceanography**. 56(4): 281-288.

- Padilla, D. K. & Klinger, T. 2005. Risks of aquaculture escapees: Oyster invader in marine reserves. **Journal of Shellfish Research**. 24: 332p.
- Perbiche-Neves, G., Favareto, L. R., de Oliveira Naliato, D. A., & Serafim-Júnior, M. 2010. Similaridade do micro-zooplâncton e relações com variáveis ambientais em um estuário subtropical. **Revista Brasileira de Biociências**. 8(1): 3-8.
- SETPT, 2012. **Guia Turístico do Litoral 2013**. Secretaria de Estado do Turismo Paraná Turismo, 103p.
- Silveira, R. C., Silva, F. C., Gomes, C. H. M., Ferreira, J. F., & Melo, C. M. R. 2011. Larval settlement and spat recovery rates of the oyster *Crassostrea brasiliana* (Lamarck, 1819) using different systems to induce metamorphosis. **Brazilian Journal of Biology**. 71(2): 557-562.
- Sorabella, L. C. & Luckenbach, M. W. 2003. A comparison of two oysters (*Crassostrea virginica*) stocks to determine suitability for use in oysters reef restoration in Virginia. **Journal of Shellfish Research**. 22: 355pp.
- Southworth, M. & Mann, R. 2004. Decadal scale changes in seasonal patterns of oyster recruitment in the Virginia sub estuaries of the Chesapeake Bay. **Journal of Shellfish Research**. 23: 312pp.
- Su, Z., Huang, L., Yan, Y. & Li, H. 2007. The effect of different substrates on pearl oyster *Pinctada martensii* (Dunker) larvae settlement. **Aquaculture**. 271: 377-383.
- Valentin, J. L. 2000. **Ecologia numérica: uma introdução à análise multivariada de dados ecológicos**. Rio de Janeiro: Interciência, 117pp.
- Wakamatsu, T. 1973. **A ostra de Cananéia e seu cultivo**. São Paulo, Superintendência do desenvolvimento do Litoral Paulista/Instituto Oceanográfico, USP, 141p.
- Wilson, C., Scotto, L., Scarpa, J., Volety, A., Laramore, S. & Haunert, D. 2005. Survey of water quality, oyster reproduction and oyster health status in the St. Lucie Estuary. **Journal of Shellfish Research**. 24: 157-165.
- Zanette, G. B., Ferreira, J. F., Silva, F. C., Pereira, A., Melo, C. M. R. 2009. Influence of the type of collector preparation periods on the setting rate the scallop *Nodipecten nodosus* L. in the laboratory. **Aquacultura Research**. 40: 1451-1458.

Received: April 2016

Accepted: December 2016

Published: January 2017